

Bayer Environmental Science



Before using this product read the label carefully / Lees die etiket noukeurig alvorens die produk gebruik word

BLATTANEX[®]

COCKROACH GEL / KAKKERLAK JEL

Reg. No. L7555 Act No. 36 of 1947
Namibian Reg. No. N-AR 0687

Reg. Nr. L7555 Wet Nr. 36 van 1947
Namibiese Reg. No. N-AR 0687

AN INSECTICIDAL GEL BAIT FOR THE
CONTROL OF COCKROACHES IN
HOMES.

'N INSEKDODENDE JEL LOKAAS VIR
DIE BEHEER VAN KAKKERLAKKE IN
HUISE.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

Imidacloprid (chloro-nicotinyl)

21,5 g/kg

Imidaklopried (chloro-nikotiniel)

NET CONTENTS / NETTO INHOUD: 5 g syringes

REGISTRATION HOLDER / REGISTRASIEHOUER:

Bayer (Pty.) Ltd. / (Edms.) Bpk.
Reg. No. / Nr. 1968/011192/07
P.O. Box / Posbus 143
ISANDO 1600
SOUTH AFRICA / SUID-AFRIKA
TEL: (011) 921 5911

Batch no: / Lotnommer:

Date of manufacture: / Datum vervaardig:

BLATTANEX[®] is a registered trademark of Bayer / is 'n geregistreerde handelsmerk van Bayer.

UN No. / VN Nr. 3077



WARNINGS:

Handle with care.
Harmful when swallowed.
Harmful to fish and other animals.
Store in a cool place and away from food and feedstuffs.
Keep out of the reach of children, uninformed persons and animals.

PRECAUTIONS:

Avoid contact by mouth or with skin.
Wash with soap and water immediately after use or accidental skin contact.
Do not eat, drink or smoke whilst applying or before washing hands and face.
Apply only in positions where it will be completely inaccessible to children and pets.
Do not contaminate dams, drains, ponds, streams and other waterways with the product or used containers.
Do not apply gel to surfaces on which food or feed is stored, prepared or eaten.
Protect food preparing equipment and eating utensils from contamination during application.
Destroy empty syringe by flattening and do not re-use container for any other purpose.

DIRECTIONS FOR USE: Use only as directed

Blattanex[®] Cockroach Gel contains a cockroach insecticide with a delayed action; therefore results can be expected within a week. Cockroaches may return to their nests before the insecticide takes effect and it is possible that dead cockroaches will not be seen in areas of application.

Blattanex[®] Cockroach Gel is applied by spot-treatment in places in or around cockroach harbourages and where cockroaches are observed or hide. The gel must be applied in dark places and out of sight, out of reach of children and animals and away from foodstuffs. Do not apply **Blattanex[®] Cockroach Gel** in areas which have been sprayed with insecticide and do not spray insecticide over the gel. Additional insecticide applications may cause the bait to become repellent and thus reduce the level of control.

Directions:

Slightly bend the tip and twist of the cap. Touch the application surface area with the tip and press the plunger slightly until the required amount of gel is applied. Replace the cap after use.

Application rates:

Infestation level	Spot size	Weight	Nr. of spots / 10 m²
Low	3 mm	0,25 g	10 – 20
High	6 mm	0,5 g	10 – 20

Numerous smaller placements will provide faster control than a few larger spots. The gel will remain pliable and palatable to cockroaches as long as it is visible and uncontaminated. Where infestations are high inspect gel four weeks after treatment and replace if necessary. The life of the bait will reduce under daylight conditions.

For any product enquiry, contact Bayer Environmental Science - 011 921 5911. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

Blattanex[®] is a registered trademark of Bayer.

© Bayer (Pty) Ltd. All unauthorized reproduction of the label is forbidden.

www.bayerhome.co.za

WAARSKUWINGS:

Hanteer versigtig.
Skadelik indien ingesluk word.
Skadelik vir visse en ander diere.
Berg in 'n koel plek weg van voer en voedsel.
Hou buite bereik van kinders, oningeligte persone en diere.

VOORSORGMAATREËLS:

Verhoed kontak deur mond en vel.
Was met seep en water na gebruik en toevallige velkontak.
Moet nie eet, drink of rook tydens toediening en voordat hande en gesig gewas is nie.
Wend aan op plekke wat totaal onbereikbaar is vir kinders en huisdiere.
Moenie damme, rioolstelsels, afvoerpype, riviere, dammetjies of ander waterweë met die produk of gebruikte houers besoedel nie.
Moenie aanwend op oppervlakke waar voedsel of voer geberg, voorberei of geëet word nie.
Voorkom besoedeling van enige voedsel voorbereidings toerusting en eetgerei tydens aanwending.
Vernietig leë spuit deur dit plat te slaan en moet houer vir geen doel hergebruik nie.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

Blattanex® Kakkerlak Jel bevat 'n kakkerlak insekdoder met 'n vertraagde aksie; resultate kan verwag word binne 'n week na behandeling. Kakkerlakke kan na hul neste terugkeer voordat die insekdoder in werking tree en dit is dus moontlik dat dooie kakkerlakke nie in die omgewing van toediening waargeneem sal word nie.

Blattanex® Kakkerlak Jel word aangewend deur kolbehandeling in en rondom kakkerlak skuilplekke en waar kakkerlakke waargeneem word. Die jel moet aangewend word in donker plekke, buite sig, buite bereik van kinders en diere en weg van voedsel. Moenie **Blattanex® Kakkerlak Jel** toedien op plekke waar insekdoder gespuit is nie en moenie insekdoder oor die jel toedien nie. Addisionele insekdoder toedienings sal die aantreklikheid van die aas benadeel en sodoende die vlak van beheer verlaag.

Aanwysings:

Buig die punt van die spuit effens en draai die doppie af. Raak met die punt aan die toedienings oppervlak area en druk die spuit liggies totdat die vereiste hoeveelheid jel toegedien is. Plaas die doppie terug na gebruik.

Toedienings hoeveelhede:

Besmettingsvlak	Druppel grootte	Massa	Aantal druppels / 10 m²
Laag	3 mm	0,25 g	10 - 20
Hoog	6 mm	0,5 g	10 - 20

Talryke kleiner druppels sal vinniger beheer gee as 'n paar groter druppels. Die jel sal buigsaam en smaaklik bly vir kakkerlakke so lank as wat dit sigbaar en skoon van vuilgoed is. Waar swaar besmettings voorkom ondersoek die jel vier weke na behandeling en vervang indien nodig. Die leeftyd van die aas sal verminder onder daglig omstandighede.

Vir enige produkt navrae, skakel Bayer Environmental Science - 011 921 5911. Slegs vir vergiftigingsnoodgevallen kan Griffon Gif Inligtingsentrum ook gekontak word - 082 446 8946.

Blattanex® is 'n geregistreerde handelsmerk van Bayer.

© Bayer (Edms.) Bpk. Alle ongemagtigde kopiëring van die etiket is verbode.

www.bayerhome.co.za