



Bayer Environmental Science

Before using this product read the label carefully. / Lees die etiket noukeurig alvorens die produk gebruik word.

FINALE[®]

Liquid Rat and Mouse Bait/Vloeistof Rot- en Muis Lokaas

Reg. No. L5657 Act No. 36 of 1947
Namibian Reg. No. N-AR 0419

Reg. Nr. L5657 Wet Nr.36 van 1947
Namibiese Reg. Nr. N-AR 0419

A HIGHLY ACTIVE ANTICOAGULANT BAIT FOR THE CONTROL OF THE NORWAY RAT (*Rattus norvegicus*), ROOF RAT (*Rattus rattus*) AND THE HOUSE MOUSE (*Mus musculus*).

'N HOOGS AKTIEWE BLOEDSTOLWEERMIDDEL LOKAAS VIR DIE BEHEER VAN DIE NOORWEEGSE ROT (*Rattus norvegicus*), DAKROT (*Rattus rattus*) EN DIE HUISMUIS (*Mus musculus*).

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

*Difethialone (anticoagulant)

0,5 g/l

Difetialoon (bloedstolweermiddel)

NET CONTENTS: / NETTO INHOUD: 200 ml

REGISTRATION HOLDER / REGISTRASIEHOUER:

Bayer (Pty.) Ltd. / (Edms.) Bpk.
Reg. No. / Nr. 1968/011192/07
P.O. Box / Posbus 143
ISANDO 1600
SOUTH AFRICA / SUID-AFRIKA
TEL: (011) 921 5911

Batch no: / Lotnommer:

Date of manufacture: / Datum vervaardig:

FINALE[®] is a registered trademark of Bayer / is 'n geregistreerde handelsmerk van Bayer.

* A Lipha Product.

UN No. / VN Nr. 3082



VERY TOXIC
UITERS GIFTIG



WARNINGS:

Handle with extreme care.

Prevent access to baits by children, uninformed persons and animals.
Store under lock and key away from food and feedstuffs.
Poisonous if swallowed and upon prolonged contact with the skin.
Toxic to fish and wildlife.
In case of poisoning, call a doctor and make this label available to him.

PRECAUTIONS:

Avoid all contact by mouth.

Avoid contamination of any edible commodity, feed, food utensils and drinking water.
Set baiting stations where these will be inaccessible to children and domestic animals.
Wear rubber gloves and wash hands after handling.
Do not eat, drink or smoke whilst handling the bait.
Rinse empty container three times with a volume of water equal to a minimum of 10 % of that of the container. Destroy empty container and do not re-use it for any other purpose.
Bury remains of baiting liquid at end of treatment and burn or bury rodent bodies.

SYMPTOMS OF POISONING

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

FIRST AID TREATMENT

Remove patient from source of poisoning and keep him quiet and reassured. Take patient IMMEDIATELY to the nearest physician. Never give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN

Administer 5 mg/kg Vit. K₁ intravenously on the first day, followed by another administration of the same dose if the Quick time does not normalise a few hours after the first injection. Follow up with 5 mg/kg/day orally on the second and following days. The duration of the treatment is generally between 10 and 15 days.

In severe cases give small transfusions of fresh blood.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

For use in the home, on the farm and industrial premises.
Effective against warfarin resistant rats and mice.

Mix 1 part **Finale[®] Liquid** with 20 parts water (50 ml to 1 ℓ water) and place in dispensers at strategic positions out of reach of children and domestic animals in area to be cleared.
Place dispensers at 10 m intervals for rats and 2 m for mice.

For any product enquiry, contact Bayer Environmental Science - 011 921 5911. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

FINALE[®] is a registered trademark of Bayer.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.

www.bayeres.co.za

WAARSKUWINGS:

Hanteer uiters versigtig.

Hou buite bereik van kinders, oningeligte persone en diere.
Bewaar agter slot en grendel weg van voedsel en voer.
Giffig indien ingesluk en by langdurige velkontak.
Giffig vir visse en wild.
In geval van vergiftiging ontbied 'n dokter en stel hierdie etiket tot sy beskikking.

VOORSORGMATEËLS:

Vermyn enige kontak met die mond.

Vermyn besoedeling van enige eetbare produkte, voer, eetgerei en drinkwater.
Plaas lokaas uit buite bereik van kinders en troeteldiere.
Dra rubberhandskoene en was hande na hantering.
Moet nie eet, drink of rook terwyl lokaas hanteer word nie.
Spoel leë houers drie keer uit met 'n volume gelykstaande aan 'n minimum van 10 % van die houers.
Vernietig leë houers en moet dit vir geen ander doel gebruik nie.
Begrawe enige lokaasvloestof wat nie opgedrink is nie aan die einde van die behandeling en verbrand of begrawe dooie knaagdiers.

SIMPTOME VAN VERGIFTIGING

Bloeding van die tandvleis en neus, onderhuidse bloeding, bleekheid, bloeding in en rondom die knie- en elmbooggewrigte. Bloed in urine en ontlasting.

EERSTEHULP BEHANDELING

Verwyder pasiënt van die bron van vergiftiging en hou hom kalm en gerusgestel. Neem pasiënt ONMIDDELLIK na die naaste geneesheer. Moet nooit enigiets per mond aan 'n bewustelose persoon gee nie.

NOTA AAN GENEESHEER

Dien 5 mg/kg Vit. K₁ intraveneus toe op die eerste dag, gevolg deur 'n addisionele toediening van dieselfde dosis indien die protrombienne tyd nie normaliseer binne 'n paar uur na die eerste inspuiting nie. Volg op met 5 mg/kg/dag mondelings op die tweede en daaropvolgende dae. Die duur van die behandeling is gewoonlik tussen 10 en 15 dae.
In ernstige gevalle gee klein oortappings vars bloed.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI.

Vir gebruik in huise, op plase en industriële persele.
Doeltreffend teen warfarin bestande rotte en muise.

Meng 1 deel **Finale[®] Vloeistof** met 20 dele water (50 ml op 1 ℓ water) en plaas in houers op strategiese plekke buite bereik van kinders en diere in die gebied wat skoongemaak gaan word.
Plaas houers uit met 10 m intervalle vir rotte en 2 m vir muise.

Vir enige produk navrae, skakel Bayer Environmental Science - 011 921 5911. Slegs vir vergiftigingsnoodgevalle kan Griffon Gif Inligtingsentrum ook gekontak word - 082 446 8946.

FINALE[®] is 'n geregistreerde handelsmerk van Bayer.

© Bayer (Edms.) Bpk. Alle ongemagtigde kopiëring van die etiket is verbode.

www.bayeres.co.za