



Maxforce[®] IC

REG. NO. L6315 Act No. 36 of 1947
Namibian Reg. No. N-AR 0689

20 g

INSECTICIDE



Before using this product read the label carefully

Only for sale to and use by registered pest control operators.

An insecticidal gel bait for the control of the German cockroach in domestic premises, public hygiene and food storage premises, which is non-volatile and non-repellent.

ACTIVE INGREDIENT:

Imidacloprid (chloro-nicotinyl).....21.5 g/kg

REGISTRATION HOLDER: Bayer (Pty.) Ltd.

Reg. No. 1968/011192/07 P. O. Box 143

ISANDO 1600, SOUTH AFRICA, TEL: (011) 921 5911

MAXFORCE[®] is a
registered trademark
of the Bayer Group.

UN No. 3077



6 005317 000976

8097956B/1214



CAUTION





Maxforce[®] IC

INSECTICIDE

Reg. No. L6315, Act No. 36 of 1947
Namibian Reg. No. N-AR 0689

Only for sale to and use by registered pest control operators.

An insecticidal gel bait for the control of the German cockroach in domestic premises, public hygiene and food storage premises, which is non-volatile and non-repellent.

ACTIVE INGREDIENT:

Imidacloprid 21,5 g/kg

REGISTRATION HOLDER:

Bayer (Pty.) Ltd. Reg. No. 1968/011192/07
P.O. Box 143, ISANDO, 1600, SOUTH AFRICA
TEL: (011) 921 5911

MAXFORCE[®] is a registered trademark of the Bayer Group.

WARNINGS:

Handle with care.
Harmful when swallowed.
Harmful to fish and other animals.
Store in a cool place and away from food and feedstuffs.
Keep out of the reach of children, uninformed persons and animals.

PRECAUTIONS:

Avoid contact by mouth or with skin.
Wear rubber gloves during application.
Wash with soap and water immediately after use or accidental skin contact.
Do not eat, drink or smoke whilst applying or before washing hands and face.
Apply only in positions where it will be completely inaccessible to children and pets.
Do not contaminate dams, drains, ponds, streams and other waterways with the product or used containers.
Do not apply gel to surfaces on which food or feed is stored, prepared or eaten.
Protect food preparing equipment and eating utensils from contamination during application.
Destroy used container and do not re-use for any other purpose.

DIRECTIONS FOR USE:

 Use only as directed.

Suitable for use in commercial and industrial premises, food handling and processing establishments, restaurants, offices, store rooms, hospitals, schools, barracks, homes, hotels, shops, market places and warehouses.

Maxforce[®] IC contains a cockroach poison with a delayed action; therefore results are not immediately visible. An effect on the cockroach population can be expected within a week. Survey the area to be treated to identify centres of cockroach infestation. **Maxforce[®] IC** is applied by spot-treatment in places in or around cockroach harbourages and where cockroaches are observed or hide. The gel must be applied in dark places and out of sight.

Maxforce[®] IC is dispensed using a purpose-built caulking gun, which is used as follows: Insert the cartridge into the gun frame. Press the trigger until the plunger contracts the piston of the cartridge, but do not apply pressure. Remove the tapered nozzle and cut off the tip of the cartridge nozzle. Replace the tapered nozzle and cut off its tip at a point which allows a thread of gel 5 mm in diameter to be dispensed. **Maxforce[®] IC** can now be applied as required in and around buildings, equipment, furniture and electronic and mechanical equipment.

Application rates:

The lower application rate should be used for low and moderate infestations, or re-treatments and the higher rate for more severe infestations or first treatment. Bait application should equal approximately 0,25-0,5 g / spot and 10 to 20 spots / 10 m². A spot of approximately 3 mm diameter equates to 0,25 g of bait. A spot of approximately 6 mm diameter equates to 0,5 g of bait.

German cockroaches: One gelspot. (One gelspot 3 mm).

Numerous smaller placements will provide faster control than a few larger spots. The gel will remain pliable and palatable to cockroaches as long as it is visible; usually up to 20 weeks. Where infestations are high inspect gel four weeks after treatment and replace if necessary. The life of the bait will reduce under daylight conditions.

Maxforce[®] IC can be integrated with space or surface spray treatments but must not be contaminated with other insecticides or placed on freshly sprayed surfaces.

For any enquiry, contact Bayer - **0861 022 937 (08610BAYER)**.
For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.
www.bayer.co.za

Bayer

80907125B / 1214



Maxforce[®] IC

INSEKDODER

Reg. Nr. L6315, Wet Nr. 36 van 1947
Namibiese Reg. Nr. N-AR 0689

Slegs vir verkoop aan en gebruik deur geregistreerde plaagoperateurs.

'n Insekdodende jel lokaas vir die beheer van die Duitse kakkerlak in huishoudelike persele, publieke higiëne en voedsel opbergspersele, wat nie-vlugtig en nie-afwerend is.

AKTIEWE BESTANDEDEEL:

Imidakloprid..... 21,5 g/kg

REGISTRASIEHOUER:

Bayer (Edms.) Bpk. Reg. Nr. 1968/011192/07
Posbus 143, ISANDO, 1600, SUID-AFRIKA
TEL: (011) 921 5911

MAXFORCE[®] is 'n geregistreerde handelsmerk van die Bayer Groep.

WAARSKUWINGS:

Hanteer versigtig.
Skadelik indien ingesluk word.
Skadelik vir visse en ander diere.
Berg in 'n koel plek weg van voer en voedsel.
Hou buite bereik van kinders, oningeligte persone en diere.

VOORSORGMATREËLS:

Verhoed kontak deur mond en vel.
Dra rubberhandskoene tydens aanwending.
Was met seep en water na gebruik en toevallige velkontak.
Moet nie eet, drink of rook tydens toediening en voordat hande en gesig gewas is nie.
Wend aan op plekke wat totaal onbereikbaar is vir kinders en huisdiere.
Moenie damme, rioolstelsels, afvoerpype, riviere, dammetjies of ander waterweë met die produk of gebruikte houers besoedel nie.
Moenie aanwend op oppervlakke waar voedsel of voer geberg, voorberei of geëet word nie.
Voorkom besoedeling van enige voedsel voorbereidings toerusting en eetgerei tydens aanwending.
Vernietig leë houers. Moet vir geen doel hergebruik nie.

GEBRUIKSAANWYSINGS:

 Gebruik slegs soos aangedui.

Geskik vir die gebruik in handel- en nywerheidspersele, voedselverwerkings- en vervaardigings-instansies, restaurante, kantore, stookkamers, hospitale, skole, kasernes, huise, hotelle, skepe, markplekke en pakhuisse.

Maxforce[®] IC bevat 'n kakkerlaktgif met 'n vertraagde aksie, daarom is resultate nie onmiddellik sigbaar nie. 'n Effek op die kakkerlak bevolking kan ver wag tot binne 'n week na behandeling. Ondersoek die area wat behandel moet word en identifiseer die kakkerlak besmettingspunte. **Maxforce[®] IC** word aangewend deur kolbehandeling in en rondom kakkerlak skuilplekke. Die jel moet in donker plekke en buite sig aangewend word.

Maxforce[®] IC word aangewend met 'n meganiese toediener wat soos volg gebruik word: Plaas die patroon in die toediener raamwerk. Trek die sneller totdat die drukker kontak maak met die suier van die patroon, maar moenie druk toepas nie. Verwyder die vasgehegte spuitstuk en knip die voorste gedeelte van die patroon se punt af. Plaas die vasgehegte spuitstuk terug en sny die punt af sodat 'n gelstroom, 5 mm in deursnee uitgelaat kan word. **Maxforce[®] IC** kan nou toegedien word soos benodig word in en om geboue, toerusting, meubels, elektroniese en meganiese toerusting.

Toedienings verhouding:

Die laer toedienings skaal moet gebruik word vir lae en gemiddelde besmettings, of tydens opvolg behandelings. Die hoër toedienings skaal moet vir swaar besmettings of vir eerste behandelings gebruik word. Aas toedienings moet min of meer gelyk wees aan 0,25-0,5 g / spikkel en 10-20 spikkels / 10 m². 'n Spikkel van ongeveer 3 mm in deursnee is gelykstaande aan 0,25 g aas. 'n Spikkel van ongeveer 6 mm in deursnee is gelykstaande aan 0,5 g aas.

Duitse kakkerlakke: Een jel spikkel. (Een jel spikkel 3 mm).

Talyke kleiner uitplasinge sal vinniger beheer gee as 'n paar groter spikkels. Die jel sal buigsam en smaaklik bly vir kakkerlakke so lank as wat dit sigbaar is; gewoonlik tot 20 weke. Waar swaar besmettings voorkom ondersoek die jel vier weke na behandeling en vervang indien nodig. Die leeftyd van die aas sal verminder onder daglig omstandighede.

Maxforce[®] IC kan geïntegreer word met lug- of oppervlakspuit-behandelings, maar moet nie gekontamineer word met ander insekdoders of op vars bespuitte oppervlakke geplaas word nie.

Vir enige navrae, skakel Bayer - 0861 022 937 (08610BAYER).

Slegs vir vergiftigingsnoodgevalle kan Griffon Gif Inligtingentrum ook gekontak word - 082 446 8946.

© Bayer (Edms.) Bpk. Alle ongemagtigde kopiëring van die etiket is verbode.
www.bayer.co.za

Bayer

80907125A / 0213