



Maxforce[®]
PRIME

REG. NO. L8580 Act No. 36 of 1947
Namibian Reg. No. N-AR 1038

30 g

INSECTICIDE



FOR FULL PARTICULARS SEE ENCLOSED PAMPHLET

Before using this product read the label carefully

Only for sale to and use by registered pest control operators.

An insecticidal gel bait for the control of German cockroaches only in domestic premises, public hygiene and food storage premises. The product is non-volatile and non-repellent.

ACTIVE INGREDIENT: Imidacloprid (chloro-nicotinyl).... 21,5 g/kg

REGISTRATION HOLDER:

Bayer (Pty.) Ltd. Reg. No. 1968/011192/07
P. O. Box 143, ISANDO 1600,
SOUTH AFRICA, TEL: (011) 921 5911

MAXFORCE[®] is a registered trademark
of the Bayer Group.

UN No. 3077

80928831B / 1214



CAUTION





Maxforce[®] PRIME

INSECTICIDE

Reg. No. L8580, Act No. 36 of 1947
Namibian Reg. No. N-AR 1038

Only for sale to and use by registered pest control operators.

An insecticidal gel bait for the control of german cockroaches only in domestic premises, public hygiene and food storage premises. The product is non-volatile and non-repellent.

ACTIVE INGREDIENT:

Imidacloprid (chloro-nicotinyl)..... 21,5 g/kg

REGISTRATION HOLDER:

Bayer (Pty.) Ltd. Reg. No. 1968/011192/07
P.O. Box 143, ISANDO 1600, SOUTH AFRICA
TEL: (011) 921 5911

MAXFORCE[®] is a registered trademark of the Bayer Group.

CAUTION

WARNINGS: Handle with care. Harmful when swallowed. May cause sensitization by skin contact. Harmful to fish and other animals. Store in a cool place and away from food and feedstuffs. Keep out of the reach of children, uninformed persons and animals. The product will not control American Cockroaches effectively.

PRECAUTIONS: Avoid contact by mouth and skin. Wear rubber gloves during application. Wash with soap and water immediately after use or accidental skin contact. Do not eat, drink or smoke whilst applying and before washing hands and face. Apply only in positions where it will be completely inaccessible to children and pets. Do not contaminate dams, drains, ponds, streams and other waterways with the product or used containers. Do not apply gel to surfaces on which food or feed is stored, prepared or eaten. Protect food preparing equipment and eating utensils from contamination during application. Destroy used container and do not re-use for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

Suitable for use in commercial and industrial premises, food handling and processing establishments, restaurants, offices, store rooms, hospitals, schools, barracks, homes, hotels, ships, market places and warehouses.

Maxforce[®] Prime contains a cockroach poison with a delayed action; therefore results are not immediately visible. An effect on the cockroach population can be expected within a week. Survey the area to be treated to identify centres of cockroach infestation.

Maxforce[®] Prime is applied by spot-treatment in places in or around cockroach harbours and where cockroaches are observed or hide. The gel must be applied in dark places and out of sight.

5 g, 10 g, 20 g: Remove the cap of the cartridge tip. Apply pressure to the trigger to release droplets from the cartridge tip.

30 g, 60 g: Maxforce Prime is dispensed using a purpose-built caulking gun, which is used as follows: Insert the cartridge into the gun frame. Remove the cap of the cartridge tip. Insert a dispensing tip as supplied with the caulking gun. A plastic dispensing tip should be used in and around electrical installations and equipment. Press the trigger of the caulking gun until the plunger contacts the piston of the cartridge. The next application of pressure to the trigger should release a droplet of gel from the dispensing tip.

Maxforce[®] Prime can now be applied as required in and around buildings, equipment, furniture and electronic and mechanical equipment.

Application rates:

Bait application should equal approximately 0,25-0,5 g / spot and 10 to 20 spots / 10 m².

- A spot of approximately 3 mm diameter equates to 0,25 g of bait.

- A spot of approximately 6 mm diameter equates to 0,5 g of bait.

The lower application rate should be used for low and moderate German cockroach infestations or re-treatments. The higher rate should be used for more severe infestations or first treatment.

Numerous smaller placements will provide faster control than a few larger spots. The gel will remain palatable to cockroaches as long as it is visible; usually up to 12 weeks. Where infestations are high inspect gel four weeks after treatment and replace if necessary. The life of the bait will reduce under daylight conditions.

Maxforce[®] Prime can be integrated with space or surface spray treatments but must not be contaminated with other insecticides or placed on freshly sprayed surfaces.

For any enquiry, contact Bayer - **0861 022 937 (08610BAYER)**. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.
www.bayer.co.za

Bayer

80984618B / 1214



Maxforce[®] PRIME

INSEKDODER

Reg. Nr. L8580, Wet Nr. 36 van 1947
Namibiese Reg. Nr. N-AR 1038

Only for sale to and use by registered pest control operators.

'n Insekdodende jel lokaas vir die beheer van slegs duitse kakkerlakke in huishoudelike persele, publieke higiëne en voedsel opbergpersele, wat nie-vlugtig en nie-afwerend is.

AKTIEWE BESTANDEEL:

Imidakloprid (chloro-nikotinie)..... 21,5 g/kg

REGISTRASIEHOUER:

Bayer (Edms.) Bpk. Reg. Nr. 1968/011192/07

Posbus 143, ISANDO 1600

SUID-AFRIKA

TEL: (011) 921 5911

MAXFORCE[®] is 'n geregistreerde handelsmerk van die Bayer Groep.

VERSIGIG

WAARSKUWINGS: Hanteer versigtig. Skadelik indien ingesluk word. Kan vel sensitiseer deur velkontak. Skadelik vir visse en ander diere. Berg in 'n koel plek weg van voer en voedsel. Hou buite bereik van kinders, oningeligte persone en diere. Die produk sal nie Amerikaanse Kakkerlakke effektief beheer nie.

VOORSORGMATREËLS: Verhoed kontak deur mond en vel. Dra rubberhandskoene tydens aanwending. Was met seep en water na gebruik of toevallige velkontak. Moet nie eet, drink of rook tydens toediening en voordat hande en gesig gewas is nie. Wend aan op plekke wat totaal onbereikbaar is vir kinders en troeteldiere. Moenie damme, riootstelsels, afvoerpype, riviere, dammetjies of ander waterweë met die produk of gebruikte houers besoedel nie. Moenie aanwend op oppervlakke waar voedsel of voer geberg, voorberei of geëet word nie. Voorkom besoedeling van enige voedsel voorbereidings toerusting en eetgerei tydens aanwending. Vernietig leë houer. Moet vir geen doel hergebruik nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI.

Gesik vir die gebruik in handel- en nywerheidspersele, voedselhanterings- en voedselverwerkings-instansies, restaurante, kantore, stookkamers, hospitale, skole, kasernes, huise, hotelle, skepe, markplekke en pakhuisse.

Maxforce[®] Prime bevat 'n kakkerlagif met 'n vertraagde aksie, daarom is resultate nie onmiddelik sigbaar nie. 'n Effek op die kakkerlak bevolking kan verwag word binne 'n week na behandeling. Ondersoek die area wat behandel moet word en identifiseer die kakkerlak besmettingspunte.

Maxforce[®] Prime word aangewend deur kolbehandeling in en rondom kakkerlak skuilplekke en waar kakkerlakke opgemerk word. Die jel moet in donker plekke en buite sig aangewend word.

5 g, 10 g, 20 g: Verwyder die doppie van die spuitpunt. Pas druk op die sneller toe om druppels deur die spuitpunt vry te stel.

30 g, 60 g: Maxforce Prime word aangewend met 'n meganiese toediener wat soos volg gebruik word: Plaas die spuit in die toediener raamwerk. Verwyder die doppie van die spuitpunt. Plaas 'n toedieningspunt wat saam met die toediener voorsien is, in die spuitpunt. 'n Plastiese toedieningspunt moet in en rondom elektriese installasies en toerusting gebruik word. Trek die sneller van die meganiese toediener totdat die drukker kontak maak met die suier van die spuit. Tydens die volgende uitoefening van druk op die sneller behoort 'n druppel deur die toedieningspunt vrygestel te word.

Maxforce[®] Prime kan nou toegedien word soos benodig word in en rondom geboue, toerusting, meubels en elektroniese en meganiese toerusting.

Toediening hoeveelhede:

Aas toedienings moet min of meer gelyk wees aan 0,25-0,5 g / druppel en 10-20 druppels / 10 m².

- 'n Druppel van ongeveer 3 mm in deursnee is gelykstaande aan 0,25 g aas.

- 'n Druppel van ongeveer 6 mm in deursnee is gelykstaande aan 0,5 g aas.

Die laer toedienings skaal moet gebruik word vir lae en gemiddelde Duitse kakkerlak besmettings, of tydens opvolg behandelings. Die hoër toedienings skaal moet vir swaar besmettings of vir eerste behandelings gebruik word.

Talryke kleiner uitplasinge sal vinniger beheer gee as 'n paar groter druppels. Die jel sal buigsaam en smaaklik bly vir kakkerlakke so lank as wat dit sigbaar is; gewoonlik tot 12 weke. Waar swaar besmettings voorkom ondersoek die jel vier weke na behandeling en vervang indien nodig. Die leeftyd van die aas sal verminder onder daglig omstandighede.

Maxforce[®] Prime kan geïntegreer word met lug- of oppervlakspuit-behandelings, maar moet nie gekontamineer word met ander insekdoders of op vas bespuite oppervlakke geplaas word nie.

Vir enige navrae, skakel Bayer - **0861 022 937 (08610BAYER)**. Slegs vir vergiftigingsnoodgevallen kan Griffon Gif Inligtingsentrum ook gekontak word - 082 446 8946.

© Bayer (Edms.) Bpk. Alle ongemagtigde kopiëring van die etiket is verbode.

www.bayer.co.za

Bayer

80984618B / 1214