



# Racumin<sup>®</sup>

## PASTE/PASTA

## RODENTICIDE

REG. NO. L6401 Act No. 36 of 1947

Before using this product read the label carefully.

**Racumin Paste is a universal bait for rat control in all areas, including dumps and animal stables. Due to its special formulation Racumin Paste presents the following benefits: excellent palatability, stays attractive and effective, convenient handling, ready-to-use, biodegradable packaging, contains bitrex.**

#### ACTIVE INGREDIENT:

Coumatetralyl (anticoagulant)..... 0,375 g/kg

#### REGISTRATION HOLDER:

Bayer (Pty.) Ltd. Reg. No. 1968/011192/07  
P.O. Box 143, ISANDO 1600, SOUTH AFRICA  
TEL: (011) 921 5911

RACUMIN<sup>®</sup> is a registered trademark of the Bayer Group.

# 5 kg (500 x 10 g Sachets)



ZA79434146C - ARTICLE 79429924

Rats • Rotte



CAUTION  
VERSIGTIG





# Racumin<sup>®</sup>

## PASTE/PASTA

### **WARNINGS:**

Handle with care. Poisonous if swallowed. Toxic to fish, wildlife, pigs and other domestic animals. Store away from food and feedstuffs. Keep out of reach of children, uninformed persons, animals, poultry and livestock. Keep baiting material in original packaging in a cool and safe place. IN CASE OF POISONING, CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM.

### **PRECAUTIONS:**

Store and apply **Racumin<sup>®</sup> Paste** away from food and feedstuffs. After work and before meals wash hands and exposed skin. Do not smoke, eat or drink while working with the bait. Remove all remains of bait and bait packaging after treatment as well as all dead rodents and dispose of in a responsible manner. Preferably place **Racumin<sup>®</sup> Paste** in bait stations in order to limit access to rodents and out of reach of children and other animals.

### **SYMPTOMS OF HUMAN POISONING:**

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around the elbow and knee joints and blood in urine and faeces.

### **FIRST AID TREATMENT:**

If accidentally swallowed induce vomiting by tickling the back of the throat. Repeat until vomit is clear.

### **NOTE TO PHYSICIAN:**

Intravenous Vitamin K1, 50-150 mg on first day depending on prothrombin time. In severe cases small transfusions of fresh blood.

### **DIRECTIONS FOR USE:** USE ONLY AS DIRECTED.

Place the paste bait in places frequented by rats, e.g. rat holes, runways and feeding places. **The paper sachets are ready to use, do not open them.** Put down up to 200 g of paste bait at each baiting point depending on infestation level. Continue putting the bait down until no more of it is taken.

**NOTE:** Coumatetralyl, like other Coumarin compounds has a cumulative action, i.e. small quantities must be consumed on several consecutive days until the specific symptoms of poisoning appear. Consequently, it is the repeated intake of small dosages and not a single intake of a large quantity that causes eradication of the rodents. This multiple feed efficacy reduces the risk of secondary poisoning of non-target animals.

For any enquiry, contact Bayer - **0861 022 937 (08610BAYER)**. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.

Batch no: / Lotnommer: See on pack

Date of manufacture: / Datum vervaardig: Sien op houer

ZA79382871C - ARTICLE 79429924

# Bayer

### **WAARSKUWINGS:**

Hanteer versigtig. Giftig wanneer ingeneem. Giftig vir vis, wild, varke en huisdiere. Bewaar weg van voedsel en voer. Hou buite bereik van kinders, oningeligte persone, diere, pluimvee en lewende hawe. Hou lokaas materiaal in oorspronklike verpakking en bewaar in 'n koel veilige plek. ONTBIED DIE GENEESHEER EN TOON HIERDIE ETIKET AAN HOM IN GEVAL VAN VERGIFTIGING.

### **VOORSORGMATREËLS:**

Bewaar en dien **Racumin<sup>®</sup> Pasta** toe weg van voedsel en voer. Was hande en blootgestelde liggaamsdele na gebruik en voor etes. Moet nie eet, rook of drink terwyl met die lokaas gewerk word nie. Verwyder alle oorblywende lokaas en lokaasverpakking na behandeling asook alle dooie knaagdiere en raak daarvan ontslae op 'n verantwoordelike wyse. **Racumin<sup>®</sup> Pasta** moet verkieslik in lokaasstasies geplaas word om sodoeende toegang tot rotte te beperk. Dit moet so geplaas word dat kinders en ander diere dit nie kan bereik nie.

### **SIMPROME VAN MENSLIKE VERGIFTIGING:**

Bloeding van die tandvleis en neus, bloeding onder die vel, bleekheid, bloeding in en rondom the elmboog- en kniegewrigte, bloed in die uriene en ontlasting.

### **NOODHULPBEHANDELING:**

Indien per mond ingeneem, veroorsaak vomering deur die agterste gedeelte van die keel te prikkel met 'n vinger. Herhaal tot die vomeersel helder is.

### **NOTA AAN GENEESHEER:**

Binne-aarse Vitamien K1, 50-150 mg op die eerste dag afhangede van protrombientyd. In ernstige gevalle klein bloedtoertappings met vars bloed.

### **GEBRUIKSAANWYSINGS:** GEBRUIK SLEGS SOOS AANGEDUI.

Plaas pasta lokaas tesame met die papieromhulsel, in plekke wat gereeld deur rotte besoek word by rotgate, -paadjies en -eetplekke. Die papiersakkies is gereed-vir-gebruik, moet dit nie oopmaak nie. Plaas tot 200 g van die pasta lokaas by al die lokaas punte, na gelang van die infestasiestadium. Hou aan om die lokaas neer te sit tot geen aas meer geneem word nie.

**NOTA:** Kumatetralil soos ander Kumarienderivate, het 'n opbouende werking m.a.w. klein hoeveelhede moet op opeenvolgende dae ingeneem word totdat die spesifieke simptome van vergiftiging voorkom. Dit is gevolglik die herhaaldeike inname van klein hoeveelhede en nie 'n enkele inname van 'n groot dosis nie, wat tot gevolg het dat knaagdiere uitgeroei word. Hierdie veelvuldige voedingseffek beperk die risiko van sekondêre vergiftiging van nie-teikendiere tot 'n minimum.

Vir enige navrae, skakel Bayer - **0861 022 937 (08610BAYER)**. Slegs vir vergiftigingsnoodgevallen kan Griffon Gif Inligtingsentrum ook gekontak word - 082 446 8946.

© Bayer (Edms.) Bpk. Alle ongemagtigde kopiëring van die etiket is verbode.  
www.bayer.co.za