



Racumin®

TRACKING POWDER

RODENTICIDE

REG. NO. L2800 Act No. 36 of 1947

Before using this product read the label carefully.

An anti-coagulant poison for control of the **NORWAY RAT, ROOF RAT and HOUSE MOUSE** in and around human and animal dwellings, factories, warehouses and other storage premises.

ACTIVE INGREDIENT:
Coumatetralyl (anticoagulant).....7,5 g/kg

REGISTRATION HOLDER:
Bayer (Pty.) Ltd.
Reg. No. 1968/011192/07
P.O. Box 143, ISANDO 1600
SOUTH AFRICA
TEL: (011) 921 5911

RACUMIN® is a registered trademark of the Bayer Group.


Batch no: See container
Date of manufacture: See container

2,5 kg

80937741B / 1214



**CAUTION
VERSIGTIG**



WARNINGS:

Handle with care.
Poisonous when swallowed or inhaled.
Toxic to fish and wildlife, pigs and other domestic animals.
Store under lock and key away from food and feedstuffs.
Keep out of reach of children, uninformed persons, animals, poultry and livestock.
IN CASE OF POISONING, CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM.

PRECAUTIONS:

Do not inhale dust.
Wear rubber gloves when handling powder or bait.
Wash with soap and water immediately after accidental skin contact.
Do not eat, drink or smoke whilst mixing and applying or before washing hands and face.
Clean or destroy all mixing utensils and dispose of wash water where it will not contaminate crops, grazing, rivers and dams.
Destroy empty container and never use for any other purpose.
Prevent contamination of edible commodities, drinking water or eating utensils and areas not under treatment.
The bait or powder should be set out in such places where it will be completely inaccessible to children and domestic animals. The use of bait boxes or other special containers is strongly recommended.
Unconsumed bait residues must be removed and burned or buried at the end of treatment.

SYMPTOMS OF HUMAN POISONING:

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around the elbow and knee joints, blood in urine and faeces.

FIRST AID TREATMENT:

Skin contact - immediately remove contaminated clothing.
Wash affected area with plenty of water and soap for at least 20 minutes. If accidentally swallowed call a physician immediately.

NOTE TO PHYSICIAN:

Intravenous Vitamin K₁, 50-150 mg on first day depending on prothrombin time. In severe cases small transfusions of fresh blood.

For any enquiry, contact Bayer - **0861 022 937 (08610BAYER)**. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.

www.bayer.co.za

Bayer

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

| PEST | DOSAGE | REMARKS |
|----------------------|---|---|
| Norway Rat, Roof Rat | Undiluted One part + 15-20 parts bait material | Tracking Powder: The Racumin® Tracking Powder should be sprinkled, not too thinly, in rat holes, on rat runs and around hiding places. Leave the powder down for at least 5 days and replenish as required. Bait: Thoroughly mix one part by mass of Racumin® Tracking Powder with 15-20 parts of bait material e.g. 30 g in 450-600 g bait material and set out in places frequented by rats. Very good effect is obtained with dry baiting material made up as follows: 60 % crushed maize, 35 % rolled oats, 5 % sugar. Traces of finely cut raw bacon. The bait must be laid out on at least 5 successive days or as long as the rats consume it. It is essential that the bait remains in a good, attractive condition during this period. |
| House Mouse | As above | Apply the powder and poisoned baiting material in the same way as for rats. Racumin® is strongly recommended for joint control of mice and rats or when it is uncertain from damage caused whether the area is infested by rats or mice. |

NOTE:

Coumatetralyl, like other Coumarin compounds has a cumulative action, i.e. small quantities must be consumed on several consecutive days until the specific symptoms of poisoning appear. Consequently, it is the repeated intake of small dosages and not a single intake of a large quantity that causes eradication of the rodents. This multiple feed efficacy reduces the risk of secondary poisoning of non-target animals.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI.

| PLAAG | TOEDIENING | OPMERKINGS |
|------------------------|---|--|
| Noorweegse Rot, Dakrot | Onverdun Een deel + 15-20 dele aas materiaal | Poeier vir bestrooiing van looppaadjies: Die Racumin® Spoorpoeier moet vrylik in rotgate, op rotpaadjies en om die wegkruipplekke van rotte gestrooi word. Laat die poeier vir ten minste vyf dae en vul aan soos benodig. Lokaas: Meng volgens massa een deel Racumin® Spoorpoeier met 15-20 dele lokaasmateriaal, bv. 30 g in 450-600 g lokaasmateriaal en plaas die mengsel waar rotte dikwels opgemerk word. Goeie resultate word verkry met droë lokaasmateriaal wat soos volg saamgestel is: 60 % gebreekte mielies, 35 % gerolde hawer, 5 % suiker. 'n Paar klein stukkie gesnyde rou spek. Die lokaas moet vir ten minste 5 opeenvolgende dae uitgesit word - of so lank as wat die rotte daarvan eet. Dit is belangrik dat die lokaas in 'n goeie, aantrekklike toestand bly gedurende hierdie tydperk. |
| Huis muis | Soos hierbo | Dien die poeier en vergiftigde lokaas op dieselfde manier toe as vir rotte. Racumin® word sterk aanbeveel vir die gelyktydige beheer van muis en rotte of wanneer daar nie definitief vasgestel kan word of die skade deur rotte of muis aangerig word nie. |

NOTA:

Koumatetralil soos ander Kumarienderivate, het 'n opbouende werking m.a.w. klein hoeveelhede moet op opeenvolgende dae ingeneem word, totdat die spesifieke simptome van vergiftiging voorkom. Dit is gevolglik die herhaaldelike inname van klein hoeveelhede en nie 'n enkele inname van 'n groot dosis nie, wat tot gevolg het dat knaagdiere uitgeroei word. Hierdie veelvuldige voedingseffek beperk die risiko van sekondêre vergiftiging van nie-teikendiere tot 'n minimum.

WAARSKUWINGS:

Hanteer versigtig.
Giftig wanneer ingeneem of ingesem.
Giftig vir vis, wild, varke en huisdiere.
Bewaar agter slot en grendel weg van voedsel en voer.
Hou buite bereik van kinders, oningeligte persone, diere, pluimvee en lewendende hawe.
ONTBIED DIE GENEESHEER EN TOON HIERDIE ETIKET AAN HOM IN GEVAL VAN VERGIFTIGING.

VOORSORGMATREËLS:

Moenie die poeier inasem nie.
Dra rubberhandskoene wanneer die poeier of lokaas hanteer word.
Was dadelik met seep en water na gebruik of toevallige velkontak.
Moet nie eet, drink of rook tydens vermenging, toediening en voordat hande en gesig gewas is nie.
Maak gereedskap deeglik skoon na gebruik of vernietig dit - raak van uitspoelwater ontslae waar dit nie gewasse, weiding, riviere en damme sal besoedel nie.
Vernietig leë houers. Moet vir geen doel weer gebruik nie.
Voorkom besoedeling van enige eetbare produk, drinkwater, eetgerei en areas nie onder behandeling nie.
Tref alle moontlike voorsorgmaatreëls om die lokaas of poeier onbereikbaar te maak vir kinders en diere. Die gebruik van lokaassissies of ander spesiale houers word sterk aanbeveel.
Nadat rotte en muis bestry is moet onverbruikte lokaas en poeier verwyder en verbrand of begrawe word.

SIMPOME VAN MENSLIKE VERGIFTIGING:

Bloeding van die tandvleis en neus, bloeding onder die vel, bleekheid, bloeding in en rondom die elmboog- en kniegewrigte, bloed in die urine en ontlasting.

NOODHULPBEHANDELING:

Velkontak - verwyder besoedelde klere onmiddellik en was besmette area met groot hoeveelhede water en seep vir ten minste 20 minute. Indien per mond ingeneem kontak 'n geneesheer onmiddellik.

NOTA AAN GENEESHEER:

Binne-aarse Vitamien K₁, 50-150 mg op die eerste dag afhange van protrombientyd. In ernstige gevalle klein oortappings vars bloed.

Vir enige navrae, skakel Bayer - **0861 022 937 (08610BAYER)**. Slegs vir vergiftigingsnoodgevälle kan Griffon Gift Inligtingentrum ook gekontak word - 082 446 8946.

© Bayer (Edms.) Bpk. Alle ongemagtigde kopiering van die etiket is verbode.

www.bayer.co.za



6 005317 000952

Bayer