

Bayer Environmental Science



Before using this product read the label carefully. / Lees die etiket noukeurig alvorens die produk gebruik word.

TEMPO®
Fogger/Beroker

Reg. No. L6878 Act No. 36 of 1947
Namibian Reg. No. N-AR 0799

Reg. Nr. L6878 Wet Nr. 36 van 1947
Namibiese Reg. Nr. N-AR 0799

KILLS COCKROACHES (GERMAN AND AMERICAN), TOBACCO BEETLE, SKIN AND HIDE BEETLE LARVAE, CONFUSED FLOUR BEETLE, DRIED FRUIT MOTH (ADULTS), RICE WEEVIL, FLEAS, FLIES, MOSQUITOES, BEDBUGS, ANTS AND MEALMOTHS.

DOOD KAKKERLAKKE (DITSE EN AMERIKAANSE), TABAKKEWER, HUID-EN VELKEWER LARWES, MALKOP-MEELKEWER, DROËVRUGTE MOT (VOLWASSENES), RYSKALANDER, VLOOIE, VLIË, MUSKIETE, WEELUISE, MIERE EN MEELMOTTE.

ACTIVE INGREDIENTS/AKTIEWE BESTANDELE:

Cyfluthrin (pyrethroid)	1,0 g/kg	Siflutrien (piretroïed)
Triflumuron (benzoyl phenylurea)	2,5 g/kg	Triflumuron (benzoyl phenylurea)
Transfluthrin (pyrethroid)	0,4 g/kg	Transflutrien (piretroïed)

NET CONTENTS: / NETTO INHOUD: 325 ml

REGISTRATION HOLDER / REGISTRASIEHOUER:

Bayer (Pty.) Ltd. / (Edms.) Bpk.
Reg. No. / Nr. 1968/011192/07
P.O. Box / Posbus 143
ISANDO 1600
SOUTH AFRICA / SUID-AFRIKA
TEL: (011) 921 5911

See bottom of can for Batch No. and Date of Manufacture.
Sien onderkant van kannetjie vir Lotnommer en Datum van Vervaardiging.

TEMPO® is a registered trademark of Bayer / is 'n geregistreerde handelsmerk van Bayer.

UN No. / VN Nr. 1950



**HARMFUL
SKADELIK**



Contains: Triflumuron which is a chitin inhibitor.

WARNINGS:

Handle with care.

May cause irritation of the skin, eyes, mucous membranes and skin sensitization.

Very toxic to aquatic organisms.

Keep out of reach of children and animals.

Highly flammable: Avoid naked sparks and flames.

Do not leave on or near hot surfaces or expose to temperatures exceeding 50 °C or direct sunlight.

Do not place on electrical installations.

Pressurised: Do not puncture or burn container, even when empty.

PRECAUTIONS:

Avoid skin contact and excessive inhalation.

Do not use in the presence of persons with known sensitivity to pyrethroids.

In cases of signs of sensitivity, obtain medical advice.

Wash hands with soap and water after use.

After accidental contact with skin, wash with copious amounts of water and soap.

Do not eat, drink or smoke whilst applying or before washing hands.

Avoid contamination of food, feeds, drinking water and eating utensils. Remove or cover exposed food, dishes, utensils and food processing equipment, also cover surfaces which come in direct contact with food.

Do not contaminate dams, drains, ponds, streams and other waterways with the product or used containers.

Wear respirator, goggles and protective clothing, if entering area during application.

Remove pets and cover or remove aquaria and all humans must evacuate area to be treated.

May stain plastic material and polished surfaces.

SYMPTOMS OF POISONING

Excitement, gastrointestinal discomfort, tremor, dizziness, headache, listlessness, nausea and vomiting, epigastric pain, muscular fasciculation of limbs, unconsciousness, convulsions and coma (very high doses).

FIRST AID AND TREATMENT

Remove patient from source of poisoning to a well-ventilated area and keep him quiet and reassured. Remove contaminated clothing and rinse contaminated body area thoroughly with plenty of soap and water. After eye contact, flush contamination out of eyes with clean water and call a doctor. If swallowed call a doctor immediately.

NOTE TO PHYSICIAN

Treat symptomatically.

Tempo® Fogger has been specially formulated to produce a microfine spraymist to flush out and kill insects hiding in cracks, crevices and places hard to reach.

Tempo® Fogger can be used in commercial and industrial premises, food handling and processing establishments, offices, store rooms, hospitals, schools, homes, hotels, restaurants, farms, ships, warehouses and silos.

Tempo® Fogger kills cockroaches (German and American), tobacco beetle, skin and hide beetle larvae, confused flour beetle, dried fruit moth (adults), rice weevil, fleas, flies, mosquitoes, bedbugs,

ants and mealmoths.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

DO NOT DEPRESS ACTUATOR UNTIL READY FOR USE.

Close doors and windows of the room to be treated. Switch off ventilation.

Ensure all applicable precautionary measures have been taken.

Open all cupboards, cabinets and doors to ensure access to those areas to be treated.

Use one 325 ml can per 160 m³.

Once the trigger is pressed, total content is released.



1. Place can on a table or stand, in the centre of the area to be treated. Cover surfaces close to the fogger can with layers of newspaper, or towels.



2. Point nozzle away and depress actuator until it locks in position (nozzle spray straight up). Set fogger upright and place on centre of covered stand. Immediately leave the room, closing the door. Keep treated area closed for at least three hours.



3. Ventilate the area for 30 minutes by opening doors and windows, before re-occupation.

For any product enquiry, contact Bayer Environmental Science - 011 921 5911. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

TEMPO® is a registered trademark of Bayer.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.

www.bayerhome.co.za

Bevat: Triflumuron wat 'n chitien inhibeerder is.

WAARSKUWINGS:

Hanteer versigtig.

Kan vel, oë en slymvliese irriteer en vel sensitiseer.

Uiters giftig vir waterlewende organismes.

Hou buite bereik van kinders en diere.

Hoogs ontvlambaar: Vermoed oop vlamme en vonke.

Moenie naby of op warm oppervlakke, of aan temperature wat 50 °C oorskry, of aan direkte sonlig blootstel nie.

Moenie bo-op elektriese installerings plaas nie.

Onder druk: Moenie die houers prik of verbrand nie, al is dit leeg.

VOORSORGMAATREËLS:

Voorkom aanraking met vel en oormatige inaseming.

Moet nie in die teenwoordigheid van persone wie sensitief is vir piretroïede, gebruik nie.

In die geval van tekens van sensitiwiteit, kry mediese hulp.

Was hande met seep en water na gebruik.

Indien per ongeluk in aanraking met die vel kom, moet dit deeglik afgewas word met genoegsame seep en water.

Moenie eet, drink of rook tydens toediening, of alvorens die hande gewas is nie.

Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei. Verwyder, of bedek blootgestelde voedsel, skottelgoed, eetgerei en voedselverwerkingsapparaat. Bedek ook oppervlakke wat direk met voedsel in aanraking kom.

Moenie damme, afvoerkanale, panne, waterstrome of enige waterweë met die produk of die houers besoedel nie.

Dra 'n respirator, skermbrille, en beskermingsklere wanneer 'n gebied betree word tydens toediening. Verwyder troeteldiere en bedek of verwyder vistenks en alle persone moet die area wat behandel gaan word verlaat.

Kan plastiek- en gepoleerde oppervlakke vlek.

SIMPTOME VAN VERGIFTIGING

Angstigtheid, spysverteringskanaal ongemak, rillings, duiseligheid, hoofpyn, lusteloosheid, naarheid en braking, maagpyn, spier fassikulasie van ledemate, bewusteloosheid, stuiptrekkings en koma (baie hoë dosisse).

EERSTEHULP BEHANDELING

Verwyder pasiënt van die bron van vergiftiging na 'n goed geventileerde omgewing en hou hom kalm en gerusgestel. Verwyder besoedelde klere en spoel die besmette liggaamsdele deeglik met baie seep en koue water af. Na oogkontak, spoel besmette oë uit met skoon water en kontak 'n geneesheer. Indieningesluk kontak 'n geneesheer onmiddellik.

NOTA AAN GENEESHEER

Behandel simptome.

Tempo® Fogger is spesiaal geformuleer om 'n fyn mis-sproei te vorm wat insekte uitdryf uit hul skuilplekke soos krake, skeure en ander plekke wat moeilik bereikbaar is, en dan dood tydens kontak.

Tempo® Fogger kan gebruik word in kommersiële en nywerheidsgeboue, voedselhanterings- en prosesseringsinstansies, kantore, stoorkamers, hospitale, skole, huise, hotelle, restaurante, plase,

skepe, opbergpersele en silo's.

Tempo® Fogger dood kakkerlakke (Duitse en Amerikaanse), tabakkewer, huid- en velkewerlarwes, malkopmeelkewer, droëvrugte mot (volwassenes), ryskalander, vlooië, vlieë, muskiete, weeluisse, miere en meelmotte.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI.

MOENIE DIE AKTIVEERDER DRUK VOORDAT DIT NIE GEREED IS VIR GEBRUIK NIE.

Maak alle deure en vensters van die vertrek wat behandel gaan word dig toe. Skakel ventilasie af.

Verseker dat alle toepaslike voorsorgmaatreëls getref is.

Open alle kasdeure, kabinette en deure om binnedringing tot hierdie gebiede te verseker.

Gebruik een 325 ml houër per 160 m³.

Nadat die aktiveerder gedruk is, sal die hele inhoud vrygestel word.



1. Plaas die houër op 'n tafel of staander in die middel van die gebied wat behandel moet word. Bedek oppervlakke naby die beroker met koerantpapier of handdoeke.



2. Rig die spuitkop weg van jou en druk die aktiveerder totdat dit in posisie sluit (spuitstuk spuit regop). Sit beroker regop en plaas in die middel van die bedekte area. Verlaat die vertrek onmiddellik en maak die deur toe. Hou die behandelde vertrek toe vir ten minste drie ure.



3. Ventileer die gebied vir 30 minute deur die vensters en deure wyd oop te maak voordat die vertrek binne gegaan word.

Vir enige produk navrae, skakel Bayer Environmental Science - 011 921 5911. Slegs vir vergiftigingsnoodgevalle kan Griffon Gif Inligtingsentrum ook gekontak word - 082 446 8946.

TEMPO® is 'n geregistreerde handelsmerk van Bayer.

© Bayer (Edms.) Bpk. Alle ongemagtigde kopiëring van die etiket is verbode.

www.bayerhome.co.za